

$$\int \text{ArcCot}[a x]^n dx$$

■ Reference: G&R 2.822.2, CRC 444, A&S 4.4.63

■ Derivation: Integration by parts

■ Rule:

$$\int \text{ArcCot}[a x] dx \rightarrow x \text{ArcCot}[a x] + \frac{\text{Log}[1 + a^2 x^2]}{2 a}$$

■ Program code:

```
Int[ArcCot[a_.*x_],x_Symbol] :=
  x*ArcCot[a*x] + Log[1+a^2*x^2]/(2*a) /;
FreeQ[a,x]
```

■ Derivation: Integration by parts

■ Rule: If $n \in \mathbb{Z} \wedge n > 1$, then

$$\int \text{ArcCot}[a x]^n dx \rightarrow x \text{ArcCot}[a x]^n + a n \int \frac{x \text{ArcCot}[a x]^{n-1}}{1 + a^2 x^2} dx$$

■ Program code:

```
Int[ArcCot[a_.*x_]^n_,x_Symbol] :=
  x*ArcCot[a*x]^n +
  Dist[a*n,Int[x*ArcCot[a*x]^(n-1)/(1+a^2*x^2),x]] /;
FreeQ[a,x] && IntegerQ[n] && n>1
```

$$\int x^m \operatorname{ArcCot}[a x]^n dx$$

- **Derivation:** Iterated integration by parts

- **Rule:** If $n \in \mathbb{Z} \wedge n > 0$, then

$$\int x \operatorname{ArcCot}[a x]^n dx \rightarrow \frac{\operatorname{ArcCot}[a x]^n}{2 a^2} + \frac{x^2 \operatorname{ArcCot}[a x]^n}{2} + \frac{n}{2 a} \int \operatorname{ArcCot}[a x]^{n-1} dx$$

- **Program code:**

```
Int[x_*ArcCot[a_.*x_]^n_,x_Symbol] :=
  ArcCot[a*x]^n/(2*a^2) + x^2*ArcCot[a*x]^n/2 +
  Dist[n/(2*a),Int[ArcCot[a*x]^(n-1),x]] /;
FreeQ[a,x] && IntegerQ[n] && n>0
```

- **Derivation:** Iterated integration by parts

- **Rule:** If $n \in \mathbb{Z} \wedge n > 0 \wedge m > 1$, then

$$\int x^m \operatorname{ArcCot}[a x]^n dx \rightarrow \frac{x^{m-1} \operatorname{ArcCot}[a x]^n}{a^2 (m+1)} + \frac{x^{m+1} \operatorname{ArcCot}[a x]^n}{m+1} +$$

$$\frac{n}{a (m+1)} \int x^{m-1} \operatorname{ArcCot}[a x]^{n-1} dx - \frac{m-1}{a^2 (m+1)} \int x^{m-2} \operatorname{ArcCot}[a x]^n dx$$

- **Program code:**

```
Int[x^m_*ArcCot[a_.*x_]^n_,x_Symbol] :=
  x^(m-1)*ArcCot[a*x]^n/(a^2*(m+1)) + x^(m+1)*ArcCot[a*x]^n/(m+1) +
  Dist[n/(a*(m+1)),Int[x^(m-1)*ArcCot[a*x]^(n-1),x]] -
  Dist[(m-1)/(a^2*(m+1)),Int[x^(m-2)*ArcCot[a*x]^n,x]] /;
FreeQ[a,x] && IntegerQ[n] && n>0 && RationalQ[m] && m>1
```

■ Derivation: Integration by parts

■ Rule: If $n \in \mathbb{Z} \wedge n > 1$, then

$$\int \frac{\text{ArcCot}[a x]^n}{x} dx \rightarrow 2 \text{ArcCot}[a x]^n \text{ArcCoth}\left[1 - \frac{2 I}{I - a x}\right] + 2 a n \int \frac{\text{ArcCot}[a x]^{n-1} \text{ArcCoth}\left[1 - \frac{2 I}{I - a x}\right]}{1 + a^2 x^2} dx$$

■ Program code:

```
Int[ArcCot[a_.*x_]^n_/x_,x_Symbol] :=
  2*ArcCot[a*x]^n*ArcCoth[1-2*I/(I-a*x)] +
  Dist[2*a*n,Int[ArcCot[a*x]^(n-1)*ArcCoth[1-2*I/(I-a*x)]/(1+a^2*x^2),x]] /;
FreeQ[a,x] && IntegerQ[n] && n>1
```

■ Derivation: Integration by parts

■ Rule: If $n \in \mathbb{Z} \wedge n > 0$, then

$$\int \frac{\text{ArcCot}[a x]^n}{x^2} dx \rightarrow -\frac{\text{ArcCot}[a x]^n}{x} - a n \int \frac{\text{ArcCot}[a x]^{n-1}}{x (1 + a^2 x^2)} dx$$

■ Program code:

```
Int[ArcCot[a_.*x_]^n_/x_^2,x_Symbol] :=
  -ArcCot[a*x]^n/x -
  Dist[a*n,Int[ArcCot[a*x]^(n-1)/(x*(1+a^2*x^2)),x]] /;
FreeQ[a,x] && IntegerQ[n] && n>0
```

■ Derivation: Inverted iterated integration by parts special case

■ Rule: If $n \in \mathbb{Z} \wedge n > 0$, then

$$\int \frac{\text{ArcCot}[a x]^n}{x^3} dx \rightarrow -\frac{a^2 \text{ArcCot}[a x]^n}{2} - \frac{\text{ArcCot}[a x]^n}{2 x^2} - \frac{a n}{2} \int \frac{\text{ArcCot}[a x]^{n-1}}{x^2} dx$$

■ Program code:

```
Int[ArcCot[a_.*x_]^n_/x_^3,x_Symbol] :=
  -a^2*ArcCot[a*x]^n/2 - ArcCot[a*x]^n/(2*x^2) -
  Dist[a*n/2,Int[ArcCot[a*x]^(n-1)/x^2,x]] /;
FreeQ[a,x] && IntegerQ[n] && n>0
```

■ **Derivation: Inverted iterated integration by parts**

■ **Rule:** If $n \in \mathbb{Z} \wedge n > 0 \wedge m < -3$, then

$$\int x^m \operatorname{ArcCot}[a x]^n dx \rightarrow \frac{x^{m+1} \operatorname{ArcCot}[a x]^n}{m+1} + \frac{a^2 x^{m+3} \operatorname{ArcCot}[a x]^n}{m+1} + \frac{a n}{m+1} \int x^{m+1} \operatorname{ArcCot}[a x]^{n-1} dx - \frac{a^2 (m+3)}{m+1} \int x^{m+2} \operatorname{ArcCot}[a x]^n dx$$

■ **Program code:**

```
Int[x_^m_*ArcCot[a_*x_]^n_,x_Symbol] :=
  x^(m+1)*ArcCot[a*x]^n/(m+1) + a^2*x^(m+3)*ArcCot[a*x]^n/(m+1) +
  Dist[a*n/(m+1),Int[x^(m+1)*ArcCot[a*x]^(n-1),x]] -
  Dist[a^2*(m+3)/(m+1),Int[x^(m+2)*ArcCot[a*x]^n,x]] /;
FreeQ[a,x] && IntegerQ[n] && n>0 && RationalQ[m] && m<-3
```

$$\int \frac{\operatorname{ArcCot}[a x]^n}{c + d x} dx$$

■ **Derivation:** Integration by parts

■ **Rule:** If $a^2 c^2 + d^2 = 0 \wedge n \in \mathbb{Z} \wedge n > 0$, then

$$\int \frac{\operatorname{ArcCot}[a x]^n}{c + d x} dx \rightarrow -\frac{\operatorname{ArcCot}[a x]^n \operatorname{Log}\left[\frac{2c}{c+dx}\right]}{d} - \frac{a n}{d} \int \frac{\operatorname{ArcCot}[a x]^{n-1} \operatorname{Log}\left[\frac{2c}{c+dx}\right]}{1 + a^2 x^2} dx$$

■ **Program code:**

```
Int[ArcCot[a_.*x_]^n_./(c_+d_.*x_),x_Symbol] :=
  -ArcCot[a*x]^n*Log[2*c/(c+d*x)]/d -
  Dist[a*n/d,Int[ArcCot[a*x]^(n-1)*Log[2*c/(c+d*x)]/(1+a^2*x^2),x] /;
FreeQ[{a,c,d},x] && ZeroQ[a^2*c^2+d^2] && IntegerQ[n] && n>0
```

$$\int \frac{x^m \operatorname{ArcCot}[a x]^n}{c + d x} dx$$

■ Derivation: Integration by parts

■ Rule: If $a^2 c^2 + d^2 = 0 \wedge n \in \mathbb{Z} \wedge n > 0$, then

$$\int \frac{\operatorname{ArcCot}[a x]^n}{x (c + d x)} dx \rightarrow \frac{\operatorname{ArcCot}[a x]^n \operatorname{Log}\left[2 - \frac{2c}{c+dx}\right]}{c} + \frac{a n}{c} \int \frac{\operatorname{ArcCot}[a x]^{n-1} \operatorname{Log}\left[2 - \frac{2c}{c+dx}\right]}{1 + a^2 x^2} dx$$

■ Program code:

```
Int[ArcCot[a_.*x_]^n_./(x_*(c_+d_.*x_)),x_Symbol] :=
  ArcCot[a*x]^n*Log[2-2*c/(c+d*x)]/c +
  Dist[a*n/c,Int[ArcCot[a*x]^(n-1)*Log[2-2*c/(c+d*x)]/(1+a^2*x^2),x] /;
FreeQ[{a,c,d},x] && ZeroQ[a^2*c^2+d^2] && IntegerQ[n] && n>0
```

■ Derivation: Integration by parts

■ Rule: If $a^2 c^2 + d^2 = 0 \wedge n \in \mathbb{Z} \wedge n > 0$, then

$$\int \frac{\operatorname{ArcCot}[a x]^n}{c x + d x^2} dx \rightarrow \frac{\operatorname{ArcCot}[a x]^n \operatorname{Log}\left[2 - \frac{2c}{c+dx}\right]}{c} + \frac{a n}{c} \int \frac{\operatorname{ArcCot}[a x]^{n-1} \operatorname{Log}\left[2 - \frac{2c}{c+dx}\right]}{1 + a^2 x^2} dx$$

■ Program code:

```
Int[ArcCot[a_.*x_]^n_./(c_.*x_+d_.*x_^2),x_Symbol] :=
  ArcCot[a*x]^n*Log[2-2*c/(c+d*x)]/c +
  Dist[a*n/c,Int[ArcCot[a*x]^(n-1)*Log[2-2*c/(c+d*x)]/(1+a^2*x^2),x] /;
FreeQ[{a,c,d},x] && ZeroQ[a^2*c^2+d^2] && IntegerQ[n] && n>0
```

■ **Derivation: Algebraic simplification**

■ **Basis:** $\frac{x}{c+dx} = \frac{1}{d} - \frac{c}{d(c+dx)}$

■ **Rule:** If $a^2 c^2 + d^2 = 0 \wedge m > 0 \wedge n \in \mathbb{Z} \wedge n > 0$, then

$$\int \frac{x^m \operatorname{ArcCot}[a x]^n}{c + d x} dx \rightarrow \frac{1}{d} \int x^{m-1} \operatorname{ArcCot}[a x]^n dx - \frac{c}{d} \int \frac{x^{m-1} \operatorname{ArcCot}[a x]^n}{c + d x} dx$$

■ **Program code:**

```
Int [x_^m_.*ArcCot[a_.*x_]^n_./(c_+d_.*x_),x_Symbol] :=
  Dist[1/d,Int[x^(m-1)*ArcCot[a*x]^n,x]] -
  Dist[c/d,Int[x^(m-1)*ArcCot[a*x]^n/(c+d*x),x]] /;
FreeQ[{a,c,d},x] && ZeroQ[a^2*c^2+d^2] && RationalQ[m] && m>0 && IntegerQ[n] && n>0
```

■ **Derivation: Algebraic simplification**

■ **Basis:** $\frac{1}{c+dx} = \frac{1}{c} - \frac{dx}{c(c+dx)}$

■ **Rule:** If $a^2 c^2 + d^2 = 0 \wedge m < -1 \wedge n \in \mathbb{Z} \wedge n > 0$, then

$$\int \frac{x^m \operatorname{ArcCot}[a x]^n}{c + d x} dx \rightarrow \frac{1}{c} \int x^m \operatorname{ArcCot}[a x]^n dx - \frac{d}{c} \int \frac{x^{m+1} \operatorname{ArcCot}[a x]^n}{c + d x} dx$$

■ **Program code:**

```
Int [x_^m_*ArcCot[a_.*x_]^n_./(c_+d_.*x_),x_Symbol] :=
  Dist[1/c,Int[x^m*ArcCot[a*x]^n,x]] -
  Dist[d/c,Int[x^(m+1)*ArcCot[a*x]^n/(c+d*x),x]] /;
FreeQ[{a,c,d},x] && ZeroQ[a^2*c^2+d^2] && RationalQ[m] && m<-1 && IntegerQ[n] && n>0
```

$$\int \frac{\text{ArcCot}[a x]^n}{c + d x^2} dx$$

- Derivation: Reciprocal rule for integration

- Rule: If $d = a^2 c$, then

$$\int \frac{1}{(c + d x^2) \text{ArcCot}[a x]} dx \rightarrow -\frac{\text{Log}[\text{ArcCot}[a x]]}{a c}$$

- Program code:

```
Int[1/((c_+d_.*x_^2)*ArcCot[a_.*x_]),x_Symbol] :=
  -Log[ArcCot[a*x]]/(a*c) /;
FreeQ[{a,c,d},x] && ZeroQ[d-a^2*c]
```

- Derivation: Power rule for integration

- Rule: If $d = a^2 c \wedge n \neq -1$, then

$$\int \frac{\text{ArcCot}[a x]^n}{c + d x^2} dx \rightarrow -\frac{\text{ArcCot}[a x]^{n+1}}{a c (n+1)}$$

- Program code:

```
Int[ArcCot[a_.*x_]^n_/(c_+d_.*x_^2),x_Symbol] :=
  -ArcCot[a*x]^(n+1)/(a*c*(n+1)) /;
FreeQ[{a,c,d,n},x] && ZeroQ[d-a^2*c] && NonzeroQ[n+1]
```


$$\int \frac{x^m \operatorname{ArcCot}[a x]^n}{c + d x^2} dx$$

■ **Derivation: Algebraic expansion**

■ **Basis:** $\frac{x}{1+a^2 x^2} = -\frac{I}{a(1+a^2 x^2)} - \frac{1}{a(I-ax)}$

■ **Rule:** If $d = a^2 c \wedge n > 0$, then

$$\int \frac{x \operatorname{ArcCot}[a x]^n}{c + d x^2} dx \rightarrow \frac{I \operatorname{ArcCot}[a x]^{n+1}}{d(n+1)} - \frac{1}{a c} \int \frac{\operatorname{ArcCot}[a x]^n}{I - a x} dx$$

■ **Program code:**

```
Int[x_*ArcCot[a_*x_]^n_/ (c_+d_*x_^2), x_Symbol] :=
  I*ArcCot[a*x]^(n+1)/(d*(n+1)) -
  Dist[1/(a*c), Int[ArcCot[a*x]^n/(I-a*x), x]] /;
FreeQ[{a,c,d}, x] && ZeroQ[d-a^2*c] && RationalQ[n] && n>0
```

■ **Derivation: Algebraic expansion**

■ **Basis:** $\frac{1}{x(1+a^2 x^2)} = -\frac{a I}{1+a^2 x^2} + \frac{I}{x(I+ax)}$

■ **Rule:** If $d = a^2 c \wedge n > 0$, then

$$\int \frac{\operatorname{ArcCot}[a x]^n}{x(c + d x^2)} dx \rightarrow \frac{I \operatorname{ArcCot}[a x]^{n+1}}{c(n+1)} + \frac{I}{c} \int \frac{\operatorname{ArcCot}[a x]^n}{x(I + a x)} dx$$

■ **Program code:**

```
Int[ArcCot[a_*x_]^n_/ (x_*(c_+d_*x_^2)), x_Symbol] :=
  I*ArcCot[a*x]^(n+1)/(c*(n+1)) +
  Dist[I/c, Int[ArcCot[a*x]^n/(x*(I+a*x)), x]] /;
FreeQ[{a,c,d}, x] && ZeroQ[d-a^2*c] && RationalQ[n] && n>0
```

■ **Derivation: Algebraic expansion**

■ **Basis:** $\frac{1}{x(1+a^2x^2)} = -\frac{a}{1+a^2x^2} + \frac{1}{x(1+ax)}$

■ **Rule:** If $d = a^2 c \wedge n > 0$, then

$$\int \frac{\text{ArcCot}[ax]^n}{cx + dx^3} dx \rightarrow \frac{1}{c(n+1)} \text{ArcCot}[ax]^{n+1} + \frac{1}{c} \int \frac{\text{ArcCot}[ax]^n}{x(1+ax)} dx$$

■ **Program code:**

```
Int[ArcCot[a_.*x_]^n_./(c_.*x_+d_.*x_^3),x_Symbol] :=
  1*ArcCot[a*x]^(n+1)/(c*(n+1)) +
  Dist[1/c,Int[ArcCot[a*x]^n/(x*(1+a*x)),x]] /;
FreeQ[{a,c,d},x] && ZeroQ[d-a^2*c] && RationalQ[n] && n>0
```

■ **Derivation: Algebraic expansion**

■ **Basis:** $\frac{x^2}{c+dx^2} = \frac{1}{d} - \frac{c}{d(c+dx^2)}$

■ **Rule:** If $d = a^2 c \wedge m > 1 \wedge n > 0$, then

$$\int \frac{x^m \text{ArcCot}[ax]^n}{c+dx^2} dx \rightarrow \frac{1}{d} \int x^{m-2} \text{ArcCot}[ax]^n dx - \frac{c}{d} \int \frac{x^{m-2} \text{ArcCot}[ax]^n}{c+dx^2} dx$$

■ **Program code:**

```
Int[x_^m*ArcCot[a_.*x_]^n_./(c_+d_.*x_^2),x_Symbol] :=
  Dist[1/d,Int[x^(m-2)*ArcCot[a*x]^n,x]] -
  Dist[c/d,Int[x^(m-2)*ArcCot[a*x]^n/(c+d*x^2),x]] /;
FreeQ[{a,c,d},x] && ZeroQ[d-a^2*c] && RationalQ[{m,n}] && m>1 && n>0
```

■ **Derivation: Algebraic expansion**

■ **Basis:** $\frac{1}{c+dx^2} = \frac{1}{c} - \frac{dx^2}{c(c+dx^2)}$

■ **Rule:** If $d = a^2 c \wedge m < -1 \wedge n > 0$, then

$$\int \frac{x^m \text{ArcCot}[ax]^n}{c+dx^2} dx \rightarrow \frac{1}{c} \int x^m \text{ArcCot}[ax]^n dx - \frac{d}{c} \int \frac{x^{m+2} \text{ArcCot}[ax]^n}{c+dx^2} dx$$

■ **Program code:**

```
Int[x_^m*ArcCot[a_.*x_]^n_./(c_+d_.*x_^2),x_Symbol] :=
  Dist[1/c,Int[x^m*ArcCot[a*x]^n,x]] -
  Dist[d/c,Int[x^(m+2)*ArcCot[a*x]^n/(c+d*x^2),x]] /;
FreeQ[{a,c,d},x] && ZeroQ[d-a^2*c] && RationalQ[{m,n}] && m<-1 && n>0
```

■ **Derivation: Integration by substitution**

■ **Basis:** If $m \in \mathbb{Z}$ or $a > 0$, $\frac{x^m \text{ArcCot}[a x]^n}{1+a^2 x^2} = -\frac{\text{Cot}[\text{ArcCot}[a x]]^n \text{ArcCot}[a x]^n}{a^{m+1}} \partial_x \text{ArcCot}[a x]$

■ **Rule:** If $d = a^2 c \wedge m, n \in \mathbb{Q} \wedge (n < 0 \vee n \notin \mathbb{Z}) \wedge (m \in \mathbb{Z} \vee a > 0)$, then

$$\int \frac{x^m \text{ArcCot}[a x]^n}{c + d x^2} dx \rightarrow -\frac{1}{a^{m+1} c} \text{Subst}\left[\int x^n \text{Cot}[x]^m dx, x, \text{ArcCot}[a x]\right]$$

■ **Program code:**

```
Int[x_^m_.*ArcCot[a_.*x_]^n_/(c_+d_.*x_^2),x_Symbol] :=
  -Dist[1/(a^(m+1)*c),Subst[Int[x^n*Cot[x]^m,x],x,ArcCot[a*x]]] /;
FreeQ[{a,c,d},x] && ZeroQ[d-a^2*c] && RationalQ[{m,n}] && (n<0 || Not[IntegerQ[n]]) && (IntegerQ[m]
```

■ **Derivation: Integration by substitution**

■ **Basis:** $\frac{x^m \text{ArcCot}[a x]^n}{1+a^2 x^2} = -\frac{1}{a} \left(\frac{\text{Cot}[\text{ArcCot}[a x]]}{a} \right)^m \text{ArcCot}[a x]^n \partial_x \text{ArcCot}[a x]$

■ **Rule:** If $d = a^2 c \wedge m, n \in \mathbb{Q} \wedge (n < 0 \vee n \notin \mathbb{Z}) \wedge \neg (m \in \mathbb{Z} \vee a > 0)$, then

$$\int \frac{x^m \text{ArcCot}[a x]^n}{c + d x^2} dx \rightarrow -\frac{1}{a c} \text{Subst}\left[\int x^n \left(\frac{\text{Cot}[x]}{a} \right)^m dx, x, \text{ArcCot}[a x]\right]$$

■ **Program code:**

```
Int[x_^m_.*ArcCot[a_.*x_]^n_/(c_+d_.*x_^2),x_Symbol] :=
  -Dist[1/(a*c),Subst[Int[x^n*(Cot[x]/a)^m,x],x,ArcCot[a*x]]] /;
FreeQ[{a,c,d},x] && ZeroQ[d-a^2*c] && RationalQ[{m,n}] && (n<0 || Not[IntegerQ[n]]) && Not[IntegerQ[m]
```

$$\int \frac{\text{ArcCot}[a x]^n \text{ArcCoth}[u]}{c + d x^2} dx$$

■ **Derivation:** Algebraic simplification

■ **Basis:** $\text{ArcCoth}[z] = \frac{1}{2} \text{Log}\left[1 + \frac{1}{z}\right] - \frac{1}{2} \text{Log}\left[1 - \frac{1}{z}\right]$

■ **Rule:** If $d = a^2 c \wedge n > 0 \wedge \left(u^2 = \left(1 - \frac{2I}{I+ax}\right)^2 \vee u^2 = \left(1 - \frac{2I}{I-ax}\right)^2\right)$, then

$$\int \frac{\text{ArcCot}[a x]^n \text{ArcCoth}[u]}{c + d x^2} dx \rightarrow \frac{1}{2} \int \frac{\text{ArcCot}[a x]^n \text{Log}\left[1 + \frac{1}{u}\right]}{c + d x^2} dx - \frac{1}{2} \int \frac{\text{ArcCot}[a x]^n \text{Log}\left[1 - \frac{1}{u}\right]}{c + d x^2} dx$$

■ **Program code:**

```
Int[ArcCot[a_*x_]^n_*ArcCoth[u_] / (c_+d_*x_^2), x_Symbol] :=
  Dist[1/2, Int[ArcCot[a*x]^n*Log[Regularize[1+1/u,x]] / (c+d*x^2), x] -
  Dist[1/2, Int[ArcCot[a*x]^n*Log[Regularize[1-1/u,x]] / (c+d*x^2), x] /;
FreeQ[{a,c,d},x] && ZeroQ[d-a^2*c] && RationalQ[n] && n>0 && (ZeroQ[u^2-(1-2*I/(I+a*x))^2] || ZeroQ[u^2-(1-2*I/(I-a*x))^2])
```

$$\int \frac{\text{ArcCot}[a x]^n \text{Log}[u]}{c + d x^2} dx$$

■ **Derivation: Integration by parts**

- **Rule:** If $d = a^2 c \wedge n > 0 \wedge (1-u)^2 = \left(1 - \frac{2I}{I+ax}\right)^2$, then

$$\int \frac{\text{ArcCot}[a x]^n \text{Log}[u]}{c + d x^2} dx \rightarrow \frac{I \text{ArcCot}[a x]^n \text{PolyLog}[2, 1-u]}{2 a c} + \frac{n I}{2} \int \frac{\text{ArcCot}[a x]^{n-1} \text{PolyLog}[2, 1-u]}{c + d x^2} dx$$

■ **Program code:**

```
Int[ArcCot[a_.*x_]^n_.*Log[u_] / (c_+d_.*x_^2), x_Symbol] :=
  I*ArcCot[a*x]^n*PolyLog[2,1-u] / (2*a*c) +
  Dist[n*I/2, Int[ArcCot[a*x]^(n-1)*PolyLog[2,1-u] / (c+d*x^2), x]] /;
FreeQ[{a,c,d}, x] && ZeroQ[d-a^2*c] && RationalQ[n] && n>0 && ZeroQ[(1-u)^2 - (1-2*I/(I+a*x))^2]
```

■ **Derivation: Integration by parts**

- **Rule:** If $d = a^2 c \wedge n > 0 \wedge (1-u)^2 = \left(1 - \frac{2I}{I-ax}\right)^2$, then

$$\int \frac{\text{ArcCot}[a x]^n \text{Log}[u]}{c + d x^2} dx \rightarrow -\frac{I \text{ArcCot}[a x]^n \text{PolyLog}[2, 1-u]}{2 a c} - \frac{n I}{2} \int \frac{\text{ArcCot}[a x]^{n-1} \text{PolyLog}[2, 1-u]}{c + d x^2} dx$$

■ **Program code:**

```
Int[ArcCot[a_.*x_]^n_.*Log[u_] / (c_+d_.*x_^2), x_Symbol] :=
  -I*ArcCot[a*x]^n*PolyLog[2,1-u] / (2*a*c) -
  Dist[n*I/2, Int[ArcCot[a*x]^(n-1)*PolyLog[2,1-u] / (c+d*x^2), x]] /;
FreeQ[{a,c,d}, x] && ZeroQ[d-a^2*c] && RationalQ[n] && n>0 && ZeroQ[(1-u)^2 - (1-2*I/(I-a*x))^2]
```

$$\int \frac{\text{ArcCot}[a x]^n \text{PolyLog}[p, u]}{c + d x^2} dx$$

■ Derivation: Integration by parts

■ Rule: If $d = a^2 c \wedge n > 0 \wedge u^2 = \left(1 - \frac{2I}{I+ax}\right)^2$, then

$$\int \frac{\text{ArcCot}[a x]^n \text{PolyLog}[p, u]}{c + d x^2} dx \rightarrow -\frac{I \text{ArcCot}[a x]^n \text{PolyLog}[p+1, u]}{2 a c} - \frac{n I}{2} \int \frac{\text{ArcCot}[a x]^{n-1} \text{PolyLog}[p+1, u]}{c + d x^2} dx$$

■ Program code:

```
Int[ArcCot[a_.x_]^n_.PolyLog[p_,u_]/(c_+d_.x_^2),x_Symbol] :=
  -I*ArcCot[a*x]^n*PolyLog[p+1,u]/(2*a*c) -
  Dist[n*I/2,Int[ArcCot[a*x]^(n-1)*PolyLog[p+1,u]/(c+d*x^2),x]] /;
FreeQ[{a,c,d,p},x] && ZeroQ[d-a^2*c] && RationalQ[n] && n>0 && ZeroQ[u^2-(1-2*I/(I+a*x))^2]
```

■ Derivation: Integration by parts

■ Rule: If $d = a^2 c \wedge n > 0 \wedge u^2 = \left(1 - \frac{2I}{I-ax}\right)^2$, then

$$\int \frac{\text{ArcCot}[a x]^n \text{PolyLog}[p, u]}{c + d x^2} dx \rightarrow \frac{I \text{ArcCot}[a x]^n \text{PolyLog}[p+1, u]}{2 a c} + \frac{n I}{2} \int \frac{\text{ArcCot}[a x]^{n-1} \text{PolyLog}[p+1, u]}{c + d x^2} dx$$

■ Program code:

```
Int[ArcCot[a_.x_]^n_.PolyLog[p_,u_]/(c_+d_.x_^2),x_Symbol] :=
  I*ArcCot[a*x]^n*PolyLog[p+1,u]/(2*a*c) +
  Dist[n*I/2,Int[ArcCot[a*x]^(n-1)*PolyLog[p+1,u]/(c+d*x^2),x]] /;
FreeQ[{a,c,d,p},x] && ZeroQ[d-a^2*c] && RationalQ[n] && n>0 && ZeroQ[u^2-(1-2*I/(I-a*x))^2]
```

$$\int \frac{\text{ArcTan}[a x]^m \text{ArcCot}[a x]^n}{c + d x^2} dx$$

■ **Derivation:** Integration by parts

■ **Rule:** If $d = a^2 c \wedge m, n \in \mathbb{Z} \wedge 0 < n \leq m$, then

$$\int \frac{\text{ArcTan}[a x]^m \text{ArcCot}[a x]^n}{c + d x^2} dx \rightarrow \frac{\text{ArcTan}[a x]^{m+1} \text{ArcCot}[a x]^n}{a c (m+1)} + \frac{n}{m+1} \int \frac{\text{ArcTan}[a x]^{m+1} \text{ArcCot}[a x]^{n-1}}{c + d x^2} dx$$

■ **Program code:**

```
Int[ArcTan[a_.x_]^m_.*ArcCot[a_.x_]^n_./(c_+d_.x_^2),x_Symbol] :=
  ArcTan[a*x]^(m+1)*ArcCot[a*x]^n/(a*c*(m+1)) +
  Dist[n/(m+1),Int[ArcTan[a*x]^(m+1)*ArcCot[a*x]^(n-1)/(c+d*x^2),x]] /;
FreeQ[{a,c,d},x] && ZeroQ[d-a^2*c] && IntegersQ[m,n] && 0<n<m
```

$$\int (c + d x^2)^m \operatorname{ArcCot}[a x]^n dx$$

- Rule: If $d = a^2 c \wedge c > 0$, then

$$\int \frac{\operatorname{ArcCot}[a x]}{\sqrt{c + d x^2}} dx \rightarrow -\frac{2 \operatorname{ArcCot}[a x] \operatorname{ArcTan}\left[\frac{\sqrt{1+I a x}}{\sqrt{1-I a x}}\right]}{a \sqrt{c}} - \frac{i \operatorname{PolyLog}\left[2, -\frac{i \sqrt{1+I a x}}{\sqrt{1-I a x}}\right]}{a \sqrt{c}} + \frac{i \operatorname{PolyLog}\left[2, \frac{i \sqrt{1+I a x}}{\sqrt{1-I a x}}\right]}{a \sqrt{c}}$$

- Program code:

```
Int[ArcCot[a_.x_]/Sqrt[c+d_.x_^2],x_Symbol] :=
  -2*I*ArcCot[a*x]*ArcTan[Sqrt[1+I*a*x]/Sqrt[1-I*a*x]]/(a*Sqrt[c]) -
  I*PolyLog[2,-I*Sqrt[1+I*a*x]/Sqrt[1-I*a*x]]/(a*Sqrt[c]) +
  I*PolyLog[2,I*Sqrt[1+I*a*x]/Sqrt[1-I*a*x]]/(a*Sqrt[c]) /;
FreeQ[{a,c,d},x] && ZeroQ[d-a^2*c] && PositiveQ[c]
```

- Basis: $\partial_x \frac{\sqrt{1+a^2 x^2}}{\sqrt{c+c a^2 x^2}} = 0$

- Rule: If $d = a^2 c \wedge \neg (c > 0)$, then

$$\int \frac{\operatorname{ArcCot}[a x]}{\sqrt{c + d x^2}} dx \rightarrow \frac{\sqrt{1 + a^2 x^2}}{\sqrt{c + d x^2}} \int \frac{\operatorname{ArcCot}[a x]}{\sqrt{1 + a^2 x^2}} dx$$

- Program code:

```
Int[ArcCot[a_.x_]/Sqrt[c+d_.x_^2],x_Symbol] :=
  Sqrt[1+a^2*x^2]/Sqrt[c+d*x^2]*Int[ArcCot[a*x]/Sqrt[1+a^2*x^2],x] /;
FreeQ[{a,c,d},x] && ZeroQ[d-a^2*c] && Not[PositiveQ[c]]
```

- Rule: If $d = a^2 c$, then

$$\int \frac{\operatorname{ArcCot}[a x]}{(c + d x^2)^{3/2}} dx \rightarrow -\frac{1}{a c \sqrt{c + d x^2}} + \frac{x \operatorname{ArcCot}[a x]}{c \sqrt{c + d x^2}}$$

- Program code:

```
Int[ArcCot[a_.x_]/(c+d_.x_^2)^(3/2),x_Symbol] :=
  -1/(a*c*Sqrt[c+d*x^2]) +
  x*ArcCot[a*x]/(c*Sqrt[c+d*x^2]) /;
FreeQ[{a,c,d},x] && ZeroQ[d-a^2*c]
```


- Rule: If $d = a^2 c \wedge n > 1$, then

$$\int \frac{\text{ArcCot}[a x]^n}{(c + d x^2)^{3/2}} dx \rightarrow -\frac{n \text{ArcCot}[a x]^{n-1}}{a c \sqrt{c + d x^2}} + \frac{x \text{ArcCot}[a x]^n}{c \sqrt{c + d x^2}} - n(n-1) \int \frac{\text{ArcCot}[a x]^{n-2}}{(c + d x^2)^{3/2}} dx$$

- Program code:

```
Int[ArcCot[a_.x_]^n_/(c_+d_.x_^2)^(3/2),x_Symbol] :=
  -n*ArcCot[a*x]^(n-1)/(a*c*Sqrt[c+d*x^2]) +
  x*ArcCot[a*x]^n/(c*Sqrt[c+d*x^2]) -
  Dist[n*(n-1),Int[ArcCot[a*x]^(n-2)/(c+d*x^2)^(3/2),x]] /;
FreeQ[{a,c,d},x] && ZeroQ[d-a^2*c] && RationalQ[n] && n>1
```

- Rule: If $d = a^2 c \wedge n < -1 \wedge n \neq -2$, then

$$\int \frac{\text{ArcCot}[a x]^n}{(c + d x^2)^{3/2}} dx \rightarrow$$

$$-\frac{\text{ArcCot}[a x]^{n+1}}{a c (n+1) \sqrt{c + d x^2}} + \frac{x \text{ArcCot}[a x]^{n+2}}{c (n+1) (n+2) \sqrt{c + d x^2}} - \frac{1}{(n+1) (n+2)} \int \frac{\text{ArcCot}[a x]^{n+2}}{(c + d x^2)^{3/2}} dx$$

- Program code:

```
Int[ArcCot[a_.x_]^n_/(c_+d_.x_^2)^(3/2),x_Symbol] :=
  -ArcCot[a*x]^(n+1)/(a*c*(n+1)*Sqrt[c+d*x^2]) +
  x*ArcCot[a*x]^(n+2)/(c*(n+1)*(n+2)*Sqrt[c+d*x^2]) -
  Dist[1/((n+1)*(n+2)),Int[ArcCot[a*x]^(n+2)/(c+d*x^2)^(3/2),x]] /;
FreeQ[{a,c,d},x] && ZeroQ[d-a^2*c] && RationalQ[n] && n<-1 && n≠-2
```

- Rule: If $d = a^2 c \wedge m > 0$, then

$$\int (c + d x^2)^m \text{ArcCot}[a x] dx \rightarrow$$

$$\frac{(c + d x^2)^m}{2 a m (2 m + 1)} + \frac{x (c + d x^2)^m \text{ArcCot}[a x]}{(2 m + 1)} + \frac{2 c m}{2 m + 1} \int (c + d x^2)^{m-1} \text{ArcCot}[a x] dx$$

- Program code:

```
Int[(c_+d_.x_^2)^m_.*ArcCot[a_.x_],x_Symbol] :=
  (c+d*x^2)^m/(2*a*m*(2*m+1)) +
  x*(c+d*x^2)^m*ArcCot[a*x]/(2*m+1) +
  Dist[2*c*m/(2*m+1),Int[(c+d*x^2)^(m-1)*ArcCot[a*x],x]] /;
FreeQ[{a,c,d},x] && ZeroQ[d-a^2*c] && RationalQ[m] && m>0
```

- Rule: If $d = a^2 c \wedge m < -1 \wedge m \neq -\frac{3}{2}$, then

$$\int (c + d x^2)^m \operatorname{ArcCot}[a x] dx \rightarrow -\frac{(c + d x^2)^{m+1}}{4 a c (m+1)^2} - \frac{x (c + d x^2)^{m+1} \operatorname{ArcCot}[a x]}{2 c (m+1)} + \frac{2 m + 3}{2 c (m+1)} \int (c + d x^2)^{m+1} \operatorname{ArcCot}[a x] dx$$

- Program code:

```
Int[(c+d*x^2)^m*ArcCot[a*x],x_Symbol] :=
  -(c+d*x^2)^(m+1)/(4*a*c*(m+1)^2) -
  x*(c+d*x^2)^(m+1)*ArcCot[a*x]/(2*c*(m+1)) +
  Dist[(2*m+3)/(2*c*(m+1)),Int[(c+d*x^2)^(m+1)*ArcCot[a*x],x]] /;
FreeQ[{a,c,d},x] && ZeroQ[d-a^2*c] && RationalQ[m] && m<-1 && m!=3/2
```

- Rule: If $d = a^2 c \wedge m < -1 \wedge m \neq -\frac{3}{2} \wedge n > 1$, then

$$\int (c + d x^2)^m \operatorname{ArcCot}[a x]^n dx \rightarrow -\frac{n (c + d x^2)^{m+1} \operatorname{ArcCot}[a x]^{n-1}}{4 a c (m+1)^2} - \frac{x (c + d x^2)^{m+1} \operatorname{ArcCot}[a x]^n}{2 c (m+1)} + \frac{2 m + 3}{2 c (m+1)} \int (c + d x^2)^{m+1} \operatorname{ArcCot}[a x]^n dx - \frac{n (n-1)}{4 (m+1)^2} \int (c + d x^2)^m \operatorname{ArcCot}[a x]^{n-2} dx$$

- Program code:

```
Int[(c+d*x^2)^m*ArcCot[a*x]^n,x_Symbol] :=
  -n*(c+d*x^2)^(m+1)*ArcCot[a*x]^(n-1)/(4*a*c*(m+1)^2) -
  x*(c+d*x^2)^(m+1)*ArcCot[a*x]^n/(2*c*(m+1)) +
  Dist[(2*m+3)/(2*c*(m+1)),Int[(c+d*x^2)^(m+1)*ArcCot[a*x]^n,x]] -
  Dist[n*(n-1)/(4*(m+1)^2),Int[(c+d*x^2)^m*ArcCot[a*x]^(n-2),x]] /;
FreeQ[{a,c,d},x] && ZeroQ[d-a^2*c] && RationalQ[{m,n}] && m<-1 && m!=3/2 && n>1
```

■ **Derivation: Integration by parts**

■ **Rule:** If $d = a^2 c \wedge m < -1 \wedge n < -1$, then

$$\int (c + d x^2)^m \operatorname{ArcCot}[a x]^n dx \rightarrow -\frac{(c + d x^2)^{m+1} \operatorname{ArcCot}[a x]^{n+1}}{a c (n+1)} + \frac{2 a (m+1)}{n+1} \int x (c + d x^2)^m \operatorname{ArcCot}[a x]^{n+1} dx$$

■ **Program code:**

```
Int[(c_+d_.*x_^2)^m_*ArcCot[a_.*x_]^n_,x_Symbol] :=
  -(c+d*x^2)^(m+1)*ArcCot[a*x]^(n+1)/(a*c*(n+1)) +
  Dist[2*a*(m+1)/(n+1),Int[x*(c+d*x^2)^m_*ArcCot[a*x]^(n+1),x]] /;
FreeQ[{a,c,d},x] && ZeroQ[d-a^2*c] && RationalQ[{m,n}] && m<-1 && n<-1
```

■ **Derivation: Integration by substitution**

■ **Basis:** If $m \in \mathbb{Z}$, $(1 + a^2 x^2)^m \operatorname{ArcCot}[a x]^n = -\frac{1}{a} \operatorname{Csc}[\operatorname{ArcCot}[a x]]^{2(m+1)} \operatorname{ArcCot}[a x]^n \partial_x \operatorname{ArcCot}[a x]$

■ **Rule:** If $d = a^2 c \wedge m \in \mathbb{Z} \wedge n \in \mathbb{Q} \wedge m < -1 \wedge (n < 0 \vee n \notin \mathbb{Z})$, then

$$\int (c + d x^2)^m \operatorname{ArcCot}[a x]^n dx \rightarrow -\frac{c^m}{a} \operatorname{Subst}\left[\int x^n \operatorname{Csc}[x]^{2(m+1)} dx, x, \operatorname{ArcCot}[a x]\right]$$

■ **Program code:**

```
Int[(c_+d_.*x_^2)^m_*ArcCot[a_.*x_]^n_,x_Symbol] :=
  -Dist[c^m/a,Subst[Int[x^n*Csc[x]^(2*(m+1)),x],x,ArcCot[a*x]]] /;
FreeQ[{a,c,d},x] && ZeroQ[d-a^2*c] && IntegerQ[m] && RationalQ[n] && m<-1 && (n<0 || Not[IntegerQ[n]])
```

■ **Basis:** If $d = a^2 c$, $D\left[\frac{c^{\frac{m-1}{2}} \sqrt{c+d x^2}}{\sqrt{1+a^2 x^2}}, x\right] = 0$

■ **Rule:** If $d = a^2 c \wedge m, n \in \mathbb{Q} \wedge m < -1 \wedge (n < 0 \vee n \notin \mathbb{Z}) \wedge m - \frac{1}{2} \in \mathbb{Z} \wedge \neg (c > 0)$, then

$$\int (c + d x^2)^m \operatorname{ArcCot}[a x]^n dx \rightarrow \frac{c^{m-\frac{1}{2}} \sqrt{c+d x^2}}{\sqrt{1+a^2 x^2}} \int (1+a^2 x^2)^m \operatorname{ArcCot}[a x]^n dx$$

■ **Program code:**

```
(* Int[(c_+d_.*x_^2)^m_*ArcCot[a_.*x_]^n_,x_Symbol] :=
  c^(m-1/2)*Sqrt[c+d*x^2]/Sqrt[1+a^2*x^2]*Int[(1+a^2*x^2)^m_*ArcCot[a*x]^n,x] /;
FreeQ[{a,c,d},x] && ZeroQ[d-a^2*c] && RationalQ[{m,n}] && m<-1 && (n<0 || Not[IntegerQ[n]]) && IntegerQ[m-1/2]
```

$$\int x^m (c + d x^2)^p \operatorname{ArcCot}[a x]^n dx$$

■ **Derivation:** Integration by parts

■ **Rule:** If $d = a^2 c \wedge p \in \mathbb{Q} \wedge n > 0 \wedge p \neq -1$, then

$$\int x (c + d x^2)^p \operatorname{ArcCot}[a x]^n dx \rightarrow \frac{(c + d x^2)^{p+1} \operatorname{ArcCot}[a x]^n}{2 d (p+1)} + \frac{n}{2 a (p+1)} \int (c + d x^2)^p \operatorname{ArcCot}[a x]^{n-1} dx$$

■ **Program code:**

```
Int[x*(c+d_.**x^2)^p_.*ArcCot[a_.**x_]^n_,x_Symbol] :=
  (c+d*x^2)^(p+1)*ArcCot[a*x]^n/(2*d*(p+1)) +
  Dist[n/(2*a*(p+1)),Int[(c+d*x^2)^p*ArcCot[a*x]^(n-1),x]] /;
FreeQ[{a,c,d},x] && ZeroQ[d-a^2*c] && RationalQ[{n,p}] && n>0 && p≠-1
```

■ **Rule:** If $d = a^2 c \wedge p \in \mathbb{Q}$, then

$$\int \frac{x (c + d x^2)^p}{\operatorname{ArcCot}[a x]^2} dx \rightarrow \frac{x (c + d x^2)^{p+1}}{a c \operatorname{ArcCot}[a x]} - \frac{1}{a} \int \frac{(1 + (2 p + 3) a^2 x^2) (c + d x^2)^p}{\operatorname{ArcCot}[a x]} dx$$

■ **Program code:**

```
Int[x*(c+d_.**x^2)^p_. / ArcCot[a_.**x_]^2,x_Symbol] :=
  x*(c+d*x^2)^(p+1)/(a*c*ArcCot[a*x]) -
  Dist[1/a,Int[(1+(2*p+3)*a^2*x^2)*(c+d*x^2)^p/ArcCot[a*x],x]] /;
FreeQ[{a,c,d},x] && ZeroQ[d-a^2*c] && RationalQ[p]
```

■ **Rule:** If $d = a^2 c \wedge n < -1 \wedge n \neq -2$, then

$$\int \frac{x \operatorname{ArcCot}[a x]^n}{(c + d x^2)^2} dx \rightarrow$$

$$- \frac{x \operatorname{ArcCot}[a x]^{n+1}}{a c (n+1) (c + d x^2)} - \frac{(1 - a^2 x^2) \operatorname{ArcCot}[a x]^{n+2}}{d (n+1) (n+2) (c + d x^2)} - \frac{4}{(n+1) (n+2)} \int \frac{x \operatorname{ArcCot}[a x]^{n+2}}{(c + d x^2)^2} dx$$

■ **Program code:**

```
Int[x*ArcCot[a_.**x_]^n_/(c+d_.**x^2)^2,x_Symbol] :=
  -x*ArcCot[a*x]^(n+1)/(a*c*(n+1)*(c+d*x^2)) -
  (1-a^2*x^2)*ArcCot[a*x]^(n+2)/(d*(n+1)*(n+2)*(c+d*x^2)) -
  Dist[4/((n+1)*(n+2)),Int[x*ArcCot[a*x]^(n+2)/(c+d*x^2)^2,x]] /;
FreeQ[{a,c,d},x] && ZeroQ[d-a^2*c] && RationalQ[n] && n<-1 && n≠-2
```

■ **Derivation: Integration by parts**

■ **Rule:** If $d = a^2 c \wedge m, n, 2p \in \mathbb{Z} \wedge m < -1 \wedge n > 0 \wedge m + 2p + 3 = 0$, then

$$\int x^m (c + d x^2)^p \operatorname{ArcCot}[a x]^n dx \rightarrow \frac{x^{m+1} (c + d x^2)^{p+1} \operatorname{ArcCot}[a x]^n}{c (m+1)} + \frac{a n}{m+1} \int x^{m+1} (c + d x^2)^p \operatorname{ArcCot}[a x]^{n-1} dx$$

■ **Program code:**

```
Int[x_^m*(c+d_.*x_^2)^p_.*ArcCot[a_.*x_]^n_,x_Symbol] :=
  x^(m+1)*(c+d*x^2)^(p+1)*ArcCot[a*x]^n/(c*(m+1)) +
  Dist[a*n/(m+1),Int[x^(m+1)*(c+d*x^2)^p*ArcCot[a*x]^(n-1),x]] /;
FreeQ[{a,c,d},x] && ZeroQ[d-a^2*c] && IntegersQ[m,n,2*p] && m<-1 && n>0 && ZeroQ[m+2*p+3]
```

■ **Derivation: Integration by parts**

■ **Rule:** If $d = a^2 c \wedge m, n, 2p \in \mathbb{Z} \wedge n < -1 \wedge m + 2p + 2 = 0$, then

$$\int x^m (c + d x^2)^p \operatorname{ArcCot}[a x]^n dx \rightarrow -\frac{x^m (c + d x^2)^{p+1} \operatorname{ArcCot}[a x]^{n+1}}{a c (n+1)} + \frac{m}{a (n+1)} \int x^{m-1} (c + d x^2)^p \operatorname{ArcCot}[a x]^{n+1} dx$$

■ **Program code:**

```
Int[x_^m*(c+d_.*x_^2)^p_.*ArcCot[a_.*x_]^n_,x_Symbol] :=
  -x^m*(c+d*x^2)^(p+1)*ArcCot[a*x]^(n+1)/(a*c*(n+1)) +
  Dist[m/(a*(n+1)),Int[x^(m-1)*(c+d*x^2)^p*ArcCot[a*x]^(n+1),x]] /;
FreeQ[{a,c,d},x] && ZeroQ[d-a^2*c] && IntegersQ[m,n,2*p] && n<-1 && ZeroQ[m+2*p+2]
```

■ **Derivation: Algebraic expansion**

■ **Basis:** $\frac{x^2}{c+dx^2} = \frac{1}{d} - \frac{c}{d(c+dx^2)}$

■ **Rule:** If $d = a^2 c \wedge m, n, 2p \in \mathbb{Z} \wedge m > 1 \wedge n \neq -1 \wedge p < -1$, then

$$\int x^m (c + dx^2)^p \operatorname{ArcCot}[ax]^n dx \rightarrow \frac{1}{d} \int x^{m-2} (c + dx^2)^{p+1} \operatorname{ArcCot}[ax]^n dx - \frac{c}{d} \int x^{m-2} (c + dx^2)^p \operatorname{ArcCot}[ax]^n dx$$

■ **Program code:**

```
Int[x_^m*(c_+d_.*x_^2)^p_*ArcCot[a_.*x_]^n_,x_Symbol] :=
  Dist[1/d,Int[x^(m-2)*(c+d*x^2)^(p+1)*ArcCot[a*x]^n,x]] -
  Dist[c/d,Int[x^(m-2)*(c+d*x^2)^p_*ArcCot[a*x]^n,x]] /;
FreeQ[{a,c,d},x] && ZeroQ[d-a^2*c] && IntegersQ[m,n,2*p] && m>1 && n≠-1 && p<-1
```

■ **Derivation: Algebraic expansion**

■ **Basis:** $\frac{1}{c+dx^2} = \frac{1}{c} - \frac{dx^2}{c(c+dx^2)}$

■ **Rule:** If $d = a^2 c \wedge m, n, 2p \in \mathbb{Z} \wedge m < 0 \wedge n \neq -1 \wedge p < -1$, then

$$\int x^m (c + dx^2)^p \operatorname{ArcCot}[ax]^n dx \rightarrow \frac{1}{c} \int x^m (c + dx^2)^{p+1} \operatorname{ArcCot}[ax]^n dx - \frac{d}{c} \int x^{m+2} (c + dx^2)^p \operatorname{ArcCot}[ax]^n dx$$

■ **Program code:**

```
Int[x_^m*(c_+d_.*x_^2)^p_*ArcCot[a_.*x_]^n_,x_Symbol] :=
  Dist[1/c,Int[x^m*(c+d*x^2)^(p+1)*ArcCot[a*x]^n,x]] -
  Dist[d/c,Int[x^(m+2)*(c+d*x^2)^p_*ArcCot[a*x]^n,x]] /;
FreeQ[{a,c,d},x] && ZeroQ[d-a^2*c] && IntegersQ[m,n,2*p] && m<0 && n≠-1 && p<-1
```

■ Derivation: Integration by parts

■ Rule: If $d = a^2 c \wedge m, n, 2p \in \mathbb{Z} \wedge m < -1 \wedge n > 0 \wedge m + 2p + 3 \neq 0$, then

$$\int x^m (c + d x^2)^p \operatorname{ArcCot}[a x]^n dx \rightarrow \frac{x^{m+1} (c + d x^2)^{p+1} \operatorname{ArcCot}[a x]^n}{c (m+1)} +$$

$$\frac{a n}{m+1} \int x^{m+1} (c + d x^2)^p \operatorname{ArcCot}[a x]^{n-1} dx - \frac{a^2 (m+2p+3)}{m+1} \int x^{m+2} (c + d x^2)^p \operatorname{ArcCot}[a x]^n dx$$

■ Program code:

```
Int[x_^m.*(c_+d_.*x_^2)^p_.*ArcCot[a_.*x_]^n_,x_Symbol] :=
  x^(m+1)*(c+d*x^2)^(p+1)*ArcCot[a*x]^n/(c*(m+1)) +
  Dist[a*n/(m+1),Int[x^(m+1)*(c+d*x^2)^p*ArcCot[a*x]^(n-1),x]] -
  Dist[a^2*(m+2*p+3)/(m+1),Int[x^(m+2)*(c+d*x^2)^p*ArcCot[a*x]^n,x]] /;
FreeQ[{a,c,d},x] && ZeroQ[d-a^2*c] && IntegersQ[m,n,2*p] && m<-1 && n>0 && NonzeroQ[m+2*p+3]
```

■ Derivation: Integration by parts

■ Rule: If $d = a^2 c \wedge m, n, 2p \in \mathbb{Z} \wedge n < -1 \wedge m + 2p + 2 \neq 0$, then

$$\int x^m (c + d x^2)^p \operatorname{ArcCot}[a x]^n dx \rightarrow -\frac{x^m (c + d x^2)^{p+1} \operatorname{ArcCot}[a x]^{n+1}}{a c (n+1)} +$$

$$\frac{m}{a (n+1)} \int x^{m-1} (c + d x^2)^p \operatorname{ArcCot}[a x]^{n+1} dx + \frac{a (m+2p+2)}{n+1} \int x^{m+1} (c + d x^2)^p \operatorname{ArcCot}[a x]^{n+1} dx$$

■ Program code:

```
Int[x_^m.*(c_+d_.*x_^2)^p_.*ArcCot[a_.*x_]^n_,x_Symbol] :=
  -x^m*(c+d*x^2)^(p+1)*ArcCot[a*x]^(n+1)/(a*c*(n+1)) +
  Dist[m/(a*(n+1)),Int[x^(m-1)*(c+d*x^2)^p*ArcCot[a*x]^(n+1),x]] +
  Dist[a*(m+2*p+2)/(n+1),Int[x^(m+1)*(c+d*x^2)^p*ArcCot[a*x]^(n+1),x]] /;
FreeQ[{a,c,d},x] && ZeroQ[d-a^2*c] && IntegersQ[m,n,2*p] && n<-1 && NonzeroQ[m+2*p+2]
```

■ **Derivation: Integration by substitution**

- **Basis:** If $p \in \mathbb{Z}$ and $(m \in \mathbb{Z} \text{ or } a > 0)$, $(e + f x^m) (1 + a^2 x^2)^p \text{ArcCot}[a x]^n =$

$$-\frac{1}{a^{m+1}} (e a^m + f \text{Cot}[\text{ArcCot}[a x]]^m) \text{Csc}[\text{ArcCot}[a x]]^{2(p+1)} \text{ArcCot}[a x]^n \partial_x \text{ArcCot}[a x]$$

- **Rule:** If $d = a^2 c \wedge m, n \in \mathbb{Q} \wedge p \in \mathbb{Z} \wedge p < -1 \wedge (n < 0 \vee n \notin \mathbb{Z}) \wedge (m \in \mathbb{Z} \vee a > 0)$, then

$$\int (e + f x^m) (c + d x^2)^p \text{ArcCot}[a x]^n dx \rightarrow -\frac{c^p}{a^{m+1}} \text{Subst}\left[\int x^n (e a^m + f \text{Cot}[x]^m) \text{Csc}[x]^{2(p+1)} dx, x, \text{ArcCot}[a x]\right]$$

■ **Program code:**

```
Int[(e_.+f_.**x_^m_.)*(c_+d_.**x_^2)^p_*ArcCot[a_.*x_]^n_,x_Symbol] :=
  -Dist[c^p/a^(m+1),Subst[Int[Expand[x^n*TrigReduce[Regularize[(e*a^m+f*Cot[x]^m)*Csc[x]^(2*(p+1)),x],
    FreeQ[{a,c,d,e,f},x] && ZeroQ[d-a^2*c] && RationalQ[{m,n}] && IntegerQ[p] && p<-1 && (n<0 || Not[IntegerQ[n]])],x],x,ArcCot[a*x]]] /;
```

■ **Derivation: Integration by substitution**

- **Basis:** If $p \in \mathbb{Z}$, $x^m (1 + a^2 x^2)^p \text{ArcCot}[a x]^n = -\frac{1}{a} \left(\frac{\text{Cot}[\text{ArcCot}[a x]]}{a} \right)^m \text{Csc}[\text{ArcCot}[a x]]^{2(p+1)} \text{ArcCot}[a x]^n \partial_x \text{ArcCot}[a x]$

- **Rule:** If $d = a^2 c \wedge m, n \in \mathbb{Q} \wedge p \in \mathbb{Z} \wedge p < -1 \wedge (n < 0 \vee n \notin \mathbb{Z}) \wedge \neg (m \in \mathbb{Z} \vee a > 0)$, then

$$\int x^m (c + d x^2)^p \text{ArcCot}[a x]^n dx \rightarrow -\frac{c^p}{a} \text{Subst}\left[\int x^n (\text{Cot}[x] / a)^m \text{Csc}[x]^{2(p+1)} dx, x, \text{ArcCot}[a x]\right]$$

■ **Program code:**

```
Int[x_^m_.*(c_+d_.**x_^2)^p_*ArcCot[a_.*x_]^n_,x_Symbol] :=
  -Dist[c^p/a,Subst[Int[x^n*(Cot[x]/a)^m*Csc[x]^(2*(p+1)),x],x,ArcCot[a*x]]] /;
  FreeQ[{a,c,d},x] && ZeroQ[d-a^2*c] && RationalQ[{m,n}] && IntegerQ[p] && p<-1 && (n<0 || Not[IntegerQ[n]])
```

- **Basis:** If $d = a^2 c$, $D\left[\frac{c^{p-\frac{1}{2}} \sqrt{c+d x^2}}{\sqrt{1+a^2 x^2}}, x\right] = 0$

- **Rule:** If $d = a^2 c \wedge m, n \in \mathbb{Q} \wedge p < -1 \wedge (n < 0 \vee n \notin \mathbb{Z}) \wedge p - \frac{1}{2} \in \mathbb{Z} \wedge \neg (c > 0)$, then

$$\int x^m (c + d x^2)^p \text{ArcCot}[a x]^n dx \rightarrow \frac{c^{p-\frac{1}{2}} \sqrt{c+d x^2}}{\sqrt{1+a^2 x^2}} \int x^m (1 + a^2 x^2)^p \text{ArcCot}[a x]^n dx$$

■ **Program code:**

```
(* Int[x_^m_.*(c_+d_.**x_^2)^p_*ArcCot[a_.*x_]^n_,x_Symbol] :=
  c^(p-1/2)*Sqrt[c+d*x^2]/Sqrt[1+a^2*x^2]*Int[x^m*(1+a^2*x^2)^p*ArcCot[a*x]^n,x] /;
  FreeQ[{a,c,d},x] && ZeroQ[d-a^2*c] && RationalQ[{m,n,p}] && p<-1 && (n<0 || Not[IntegerQ[n]]) && IntegerQ[p-1/2]
```


$$\int \text{ArcCot}[a + b x^n] \, dx$$

■ Reference: G&R 2.822.2, CRC 444, A&S 4.4.63

■ Derivation: Integration by parts

■ Rule:

$$\int \text{ArcCot}[a + b x] \, dx \rightarrow \frac{(a + b x) \text{ArcCot}[a + b x]}{b} + \frac{\text{Log}[1 + (a + b x)^2]}{2 b}$$

■ Program code:

```
Int[ArcCot[a_+b_.*x_],x_Symbol] :=
  (a+b*x)*ArcCot[a+b*x]/b + Log[1+(a+b*x)^2]/(2*b) /;
FreeQ[{a,b},x]
```

■ Reference: G&R 2.822.2, CRC 444, A&S 4.4.63

■ Derivation: Integration by parts

■ Rule: If $n \in \mathbb{Q}$, then

$$\int \text{ArcCot}[a + b x^n] \, dx \rightarrow x \text{ArcCot}[a + b x^n] + b n \int \frac{x^n}{1 + a^2 + 2 a b x^n + b^2 x^{2n}} \, dx$$

■ Program code:

```
Int[ArcCot[a_+b_.*x_^n_],x_Symbol] :=
  x*ArcCot[a+b*x^n] +
  Dist[b*n,Int[x^n/(1+a^2+2*a*b*x^n+b^2*x^(2*n)),x]] /;
FreeQ[{a,b},x] && RationalQ[n]
```

$$\int x^m \operatorname{ArcCot}[a + b x^n] \, dx$$

- **Derivation:** Algebraic expansion

- **Basis:** $\operatorname{ArcCot}[z] = \frac{1}{2} i \operatorname{Log}\left[1 - \frac{i}{z}\right] - \frac{1}{2} i \operatorname{Log}\left[1 + \frac{i}{z}\right]$

- **Rule:**

$$\int \frac{\operatorname{ArcCot}[a + b x^n]}{x} \, dx \rightarrow \frac{i}{2} \int \frac{\operatorname{Log}\left[1 - \frac{i}{a + b x^n}\right]}{x} \, dx - \frac{i}{2} \int \frac{\operatorname{Log}\left[1 + \frac{i}{a + b x^n}\right]}{x} \, dx$$

- **Program code:**

```
Int[ArcCot[a_+b_.*x_^n_]/x_,x_Symbol] :=
  Dist[I/2,Int[Log[1-I/(a+b*x^n)]/x,x]] -
  Dist[I/2,Int[Log[1+I/(a+b*x^n)]/x,x]] /;
FreeQ[{a,b,n},x]
```

- **Reference:** G&R 2.852, CRC 458, A&S 4.4.71

- **Derivation:** Integration by parts

- **Rule:** If $m, n \in \mathbb{Q} \wedge m+1 \neq 0 \wedge m+1 \neq n$, then

$$\int x^m \operatorname{ArcCot}[a + b x^n] \, dx \rightarrow \frac{x^{m+1} \operatorname{ArcCot}[a + b x^n]}{m+1} + \frac{b n}{m+1} \int \frac{x^{m+n}}{1 + a^2 + 2 a b x^n + b^2 x^{2n}} \, dx$$

- **Program code:**

```
Int[x_^m_.*ArcCot[a_+b_.*x_^n_],x_Symbol] :=
  x^(m+1)*ArcCot[a+b*x^n]/(m+1) +
  Dist[b*n/(m+1),Int[x^(m+n)/(1+a^2+2*a*b*x^n+b^2*x^(2*n)),x]] /;
FreeQ[{a,b},x] && RationalQ[{m,n}] && m+1!=0 && m+1!=n
```

$$\int \operatorname{ArcCot}[a + b x]^n dx$$

- **Derivation:** Integration by substitution

- **Rule:** If $n \in \mathbb{Z} \wedge n > 1$, then

$$\int \operatorname{ArcCot}[a + b x]^n dx \rightarrow \frac{1}{b} \operatorname{Subst}\left[\int \operatorname{ArcCot}[x]^n dx, x, a + b x\right]$$

- **Program code:**

```
Int[ArcCot[a_+b_.*x_]^n_,x_Symbol] :=
  Dist[1/b,Subst[Int[ArcCot[x]^n,x],x,a+b*x]] /;
FreeQ[{a,b},x] && IntegerQ[n] && n>1
```

$$\int x^m \operatorname{ArcCot}[a + b x]^n dx$$

- **Derivation:** Integration by substitution

- **Rule:** If $m, n \in \mathbb{Z} \wedge m > 0 \wedge n > 1$, then

$$\int x^m \operatorname{ArcCot}[a + b x]^n dx \rightarrow \frac{1}{b^{m+1}} \operatorname{Subst}\left[\int (x - a)^m \operatorname{ArcCot}[x]^n dx, x, a + b x\right]$$

- **Program code:**

```
Int[x_^m_.*ArcCot[a_+b_.*x_]^n_,x_Symbol] :=
  Dist[1/b^(m+1),Subst[Int[(x-a)^m*ArcCot[x]^n,x],x,a+b*x]] /;
FreeQ[{a,b},x] && IntegersQ[m,n] && m>0 && n>1
```

$$\int \frac{\text{ArcCot}[a + b x]}{c + d x^n} dx$$

■ **Derivation: Algebraic simplification**

■ **Basis:** $\text{ArcCot}[z] = \frac{1}{2} i \text{Log}\left[1 - \frac{i}{z}\right] - \frac{1}{2} i \text{Log}\left[1 + \frac{i}{z}\right]$

■ **Rule:** If $n \in \mathbb{Z} \wedge \neg (n = 2 \wedge d = b^2 c)$, then

$$\int \frac{\text{ArcCot}[b x]}{c + d x^n} dx \rightarrow \frac{i}{2} \int \frac{\text{Log}\left[1 - \frac{i}{b x}\right]}{c + d x^n} dx - \frac{i}{2} \int \frac{\text{Log}\left[1 + \frac{i}{b x}\right]}{c + d x^n} dx$$

■ **Program code:**

```
Int[ArcCot[b_.*x_]/(c_+d_.*x_^n_),x_Symbol] :=
  Dist[I/2,Int[Log[1-I/(b*x)]/(c+d*x^n),x]] -
  Dist[I/2,Int[Log[1+I/(b*x)]/(c+d*x^n),x]] /;
FreeQ[{b,c,d},x] && IntegerQ[n] && Not[n==2 && ZeroQ[d-b^2*c]]
```

■ **Derivation: Algebraic simplification**

■ **Basis:** $\text{ArcCot}[z] = \frac{1}{2} i \text{Log}\left[1 - \frac{i}{z}\right] - \frac{1}{2} i \text{Log}\left[1 + \frac{i}{z}\right]$

■ **Rule:** If $n \in \mathbb{Z} \wedge \neg (n = 1 \wedge a d - b c = 0)$, then

$$\int \frac{\text{ArcCot}[a + b x]}{c + d x^n} dx \rightarrow \frac{i}{2} \int \frac{\text{Log}\left[1 - \frac{i}{a + b x}\right]}{c + d x^n} dx - \frac{i}{2} \int \frac{\text{Log}\left[1 + \frac{i}{a + b x}\right]}{c + d x^n} dx$$

■ **Program code:**

```
Int[ArcCot[a_+b_.*x_]/(c_+d_.*x_^n_),x_Symbol] :=
  Dist[I/2,Int[Log[1-I/(a+b*x)]/(c+d*x^n),x]] -
  Dist[I/2,Int[Log[1+I/(a+b*x)]/(c+d*x^n),x]] /;
FreeQ[{a,b,c,d},x] && IntegerQ[n] && Not[n==1 && ZeroQ[a*d-b*c]]
```

$$\int u \operatorname{ArcCot} \left[\frac{c}{a + b x^n} \right]^m dx$$

- **Derivation:** Algebraic simplification

- **Basis:** $\operatorname{ArcCot}[z] = \operatorname{ArcTan}\left[\frac{1}{z}\right]$

- **Rule:**

$$\int u \operatorname{ArcCot} \left[\frac{c}{a + b x^n} \right]^m dx \rightarrow \int u \operatorname{ArcTan} \left[\frac{a}{c} + \frac{b x^n}{c} \right]^m dx$$

- **Program code:**

```
Int[u_.*ArcCot[c_./(a_.+b_.*x_^n_.)]^m_.,x_Symbol] :=
  Int[u*ArcTan[a/c+b*x^n/c]^m,x] /;
FreeQ[{a,b,c,n,m},x]
```

$$\int \frac{f[x, \text{ArcCot}[a + b x]]}{1 - (a + b x)^2} dx$$

■ **Derivation: Integration by substitution**

■ **Basis:** $\frac{f[z]}{1+z^2} = -f[\text{Cot}[\text{ArcCot}[z]]] \text{ArcCot}'[z]$

■ **Basis:** $r + s x + t x^2 = -\frac{s^2 - 4 r t}{4 t} \left(1 - \frac{(s + 2 t x)^2}{s^2 - 4 r t}\right)$

■ **Basis:** $1 + \text{Cot}[z]^2 = \text{Csc}[z]^2$

■ **Rule:**

$$\int \frac{f[x, \text{ArcCot}[a + b x]]}{1 + (a + b x)^2} dx \rightarrow -\frac{1}{b} \text{Subst}\left[\int f\left[-\frac{a}{b} + \frac{\text{Cot}[x]}{b}, x\right] dx, x, \text{ArcCot}[a + b x]\right]$$

■ **Program code:**

```
If[ShowSteps,

Int[u_*v_^n_,x_Symbol] :=
Module[{tmp=InverseFunctionOfLinear[u,x]},
ShowStep["", "Int[f[x,ArcCot[a+b*x]]/(1+(a+b*x)^2),x]",
"-Subst[Int[f[-a/b+Cot[x]/b,x],x,ArcCot[a+b*x]]/b",Hold[
Dist[-(-Discriminant[v,x]/(4*Coefficient[v,x,2]))^n/Coefficient[tmp[[1]],x,1],
Subst[Int[Regularize[SubstForInverseFunction[u,tmp,x]*Csc[x]^(2*(n+1)),x],x], x, tmp]]] /;
NotFalseQ[tmp] && Head[tmp]==ArcCot && ZeroQ[Discriminant[v,x]*tmp[[1]]^2+D[v,x]^2] /;
SimplifyFlag && QuadraticQ[v,x] && IntegerQ[n] && n<0 && NegQ[Discriminant[v,x]] && MatchQ[u,r_.*f_^

Int[u_*v_^n_,x_Symbol] :=
Module[{tmp=InverseFunctionOfLinear[u,x]},
Dist[-(-Discriminant[v,x]/(4*Coefficient[v,x,2]))^n/Coefficient[tmp[[1]],x,1],
Subst[Int[Regularize[SubstForInverseFunction[u,tmp,x]*Csc[x]^(2*(n+1)),x],x], x, tmp]] /;
NotFalseQ[tmp] && Head[tmp]==ArcCot && ZeroQ[Discriminant[v,x]*tmp[[1]]^2+D[v,x]^2] /;
QuadraticQ[v,x] && IntegerQ[n] && n<0 && NegQ[Discriminant[v,x]] && MatchQ[u,r_.*f_^w_ /; FreeQ[f,x]
```

$$\int \operatorname{ArcCot}\left[a + b f^{c+d x}\right] d x$$

■ **Derivation:** Algebraic simplification

■ **Basis:** $\operatorname{ArcCot}[z] = \frac{1}{2} i \operatorname{Log}\left[1 - \frac{i}{z}\right] - \frac{1}{2} i \operatorname{Log}\left[1 + \frac{i}{z}\right]$

■ **Rule:**

$$\int \operatorname{ArcCot}\left[a + b f^{c+d x}\right] d x \rightarrow \frac{i}{2} \int \operatorname{Log}\left[1 - \frac{i}{a + b f^{c+d x}}\right] d x - \frac{i}{2} \int \operatorname{Log}\left[1 + \frac{i}{a + b f^{c+d x}}\right] d x$$

■ **Program code:**

```
Int[ArcCot[a_.+b_.*f_^(c_.+d_.*x_)],x_Symbol] :=
  Dist[I/2,Int[Log[1-I/(a+b*f^(c+d*x))],x]] -
  Dist[I/2,Int[Log[1+I/(a+b*f^(c+d*x))],x]] /;
FreeQ[{a,b,c,d,f},x]
```


$$\int x^m \operatorname{ArcCot} \left[a + b f^{c+d x} \right] dx$$

- **Derivation:** Algebraic simplification

- **Basis:** $\operatorname{ArcCot}[z] = \frac{1}{2} i \operatorname{Log} \left[1 - \frac{i}{z} \right] - \frac{1}{2} i \operatorname{Log} \left[1 + \frac{i}{z} \right]$

- **Rule:** If $m \in \mathbb{Z} \wedge m > 0$, then

$$\int x^m \operatorname{ArcCot} \left[a + b f^{c+d x} \right] dx \rightarrow \frac{I}{2} \int x^m \operatorname{Log} \left[1 - \frac{i}{a + b f^{c+d x}} \right] dx - \frac{I}{2} \int x^m \operatorname{Log} \left[1 + \frac{i}{a + b f^{c+d x}} \right] dx$$

- **Program code:**

```
Int[x_^m_*ArcCot[a_+b_*f^(c_+d_*x_)],x_Symbol] :=
  Dist[I/2,Int[x^m*Log[1-I/(a+b*f^(c+d*x))],x]] -
  Dist[I/2,Int[x^m*Log[1+I/(a+b*f^(c+d*x))],x]] /;
FreeQ[{a,b,c,d,f},x] && IntegerQ[m] && m>0
```

$$\int v \operatorname{ArcCot}[u] \, dx$$

- Derivation: Integration by parts

- Rule: If u is free of inverse functions, then

$$\int \operatorname{ArcCot}[u] \, dx \rightarrow x \operatorname{ArcCot}[u] + \int \frac{x \partial_x u}{1 + u^2} \, dx$$

- Program code:

```
Int[ArcCot[u_],x_Symbol] :=
  x*ArcCot[u] +
  Int[Regularize[x*D[u,x]/(1+u^2),x],x] /;
InverseFunctionFreeQ[u,x]
```

- Derivation: Integration by parts

- Rule: If $m + 1 \neq 0 \wedge u$ is free of inverse functions, then

$$\int x^m \operatorname{ArcCot}[u] \, dx \rightarrow \frac{x^{m+1} \operatorname{ArcCot}[u]}{m+1} + \frac{1}{m+1} \int \frac{x^{m+1} \partial_x u}{1 + u^2} \, dx$$

- Program code:

```
Int[x_^m_.*ArcCot[u_],x_Symbol] :=
  x^(m+1)*ArcCot[u]/(m+1) +
  Dist[1/(m+1),Int[Regularize[x^(m+1)*D[u,x]/(1+u^2),x],x]] /;
FreeQ[m,x] && NonzeroQ[m+1] && InverseFunctionFreeQ[u,x] &&
  Not[FunctionOfQ[x^(m+1),u,x]] &&
  FalseQ[PowerVariableExpn[u,m+1,x]]
```

- Derivation: Integration by parts

- Rule: If u is free of inverse functions, let $w = \int v \, dx$, if w is free of inverse functions, then

$$\int v \operatorname{ArcCot}[u] \, dx \rightarrow w \operatorname{ArcCot}[u] + \int \frac{w \partial_x u}{1 + u^2} \, dx$$

- Program code:

```
Int[v_*ArcCot[u_] , x_Symbol] :=
  Module[{w=Block[{ShowSteps=False, StepCounter=None}], Int[v,x]}],
  w*ArcCot[u] +
  Int[Regularize[w*D[u,x]/(1+u^2),x],x] /;
  InverseFunctionFreeQ[w,x] /;
  InverseFunctionFreeQ[u,x] &&
  Not[MatchQ[v, x^m_. /; FreeQ[m,x]]] &&
  FalseQ[FunctionOfLinear[v*ArcCot[u],x]]
```